

Changes in the viability of *Strongyloides ransomi* larvae (Nematoda, Rhabditida) under the influence of synthetic flavourings

A. A. Boyko*, V. V. Brygadyrenko**

*Dnipropetrovsk State Agrarian-Economic University, Dnipro, Ukraine

**Oles Honchar Dnipropetrovsk National University, Dnipro, Ukraine

Article info

Received 11.12.2016

Received in revised form

14.01.2017

Accepted 19.01.2017

Dnipropetrovsk State
Agrarian-Economic University,
Voroshilov Str., 25,
Dnipro, 49027, Ukraine
Tel.: +38-097-296-42-10
E-mail:
boikoalexandra1982@gmail.com

Oles Honchar Dnipropetrovsk
National University,
Gagarin Ave., 72,
Dnipro, 49010, Ukraine
Tel.: +38-050-93-90-788
E-mail: brigad@ua.fm

Boyko, A. A., & Brygadyrenko, V. V. (2017). Changes in the viability of *Strongyloides ransomi* larvae (Nematoda, Rhabditida) under the influence of synthetic flavourings. *Regulatory Mechanisms in Biosystems*, 8(1), 36–40. doi: 10.15421/021707

One of the most common nematodes of pigs globally is *Strongyloides ransomi* Schwartz and Alicata 1930. It usually causes aggravation of physiological indicators of its hosts and damage to their immune system. Also it is a good modelling object for the evaluation of the antiparasitic activity of new antihelminthic drugs. We conducted laboratory experiments to assess the effect of flavouring additives with flower odour (benzaldehyde, citral, D-limonene and β -ionone) upon the viability of *S. ransomi* larvae. The mortality rate was calculated for 24 hours exposure at four concentrations of each substance (10, 1, 0.1 и 0.01 g/l) with eight replications. The lowest LD₅₀ values were obtained for citral (97 mg/l) and benzaldehyde (142 mg/l). These substances are recommended for further evaluation of their antihelminthic effect in experiments using laboratory animals. Unlike other substances, the effect of β -ionone and D-limonene even at a concentration of 10 g/l after 24 hours caused the death of <50% of *S. ransomi* larvae. The study of flavouring additives with flowery odour, which are permitted to be used in food for humans and also to be used in cosmetics, is a promising field for research aimed at the development of new antiparasitic drugs.

Keywords: nematodes; antiparasitic activity; flavouring agents; benzaldehyde; citral; D-limonene; β -ionone

Introduction

Helminthiasis causes significant economic damage to animal husbandry (Faye et al., 2003; Veneziano et al., 2004; Charlier et al., 2007; Cringoli, 2008; Yamov and Antropov, 2008; Ponomar et al., 2014a, b; Boyko et al., 2016). Pig farming is one of the most common sources of meat products. Among the parasites of pigs, the most common are nematodes, including strongyloidiasis agents (Knecht et al., 2011; Eysker et al., 2011; Samsonovich, 2012a; Maslova et al., 2015). Parasitisation by *Strongyloides ransomi* (Schwartz and Alicata, 1930) causes loss of concentration of albuminous and albumin fractions in the blood of piglets, loss of erythrocytes, hemoglobin, albumin, phagocytic and lysozyme activity and immune reaction (Samsonovich, 2012b).

Therefore determining the viability of helminths is significant for controlling their population, both in the host, and in the environment. Nowadays, synthetic antiparasitic drugs (Ponomar et al., 2013) are used and experiments of identifying the antihelminthic properties of plants are being conducted (Rahmann and Seip, 2006; Burke et al., 2009; Lu et al., 2010).

To fight pathogenic organisms, microbiologists and virusologists are researching the effect of flavouring agents in food production (Chiang et al., 2005; Sato et al., 2006; Somolinos et al., 2008; Si et al., 2009; Belletti et al., 2010). Because pigs consume food not only of vegetable origin, but also of animal origin, their diet can include synthetic flavouring agents from the diet of humans. Therefore the objective of this research is to define the effect of flavouring agents upon the level of viability of *S. ransomi* larvae (Rhabditida, Strongyloididae).

Materials and methods

The faeces of pigs were studied to find *S. ransomi* larvae using the Baermann test (Zajac et al., 2011). The material was collected in the summer period in Dnipropetrovsk district, and then transferred to the laboratory in plastic containers at a temperature of 22–24 °C. *S. ransomi* culture is represented by larval stages of freely moving and parasitic forms (Van Wyk and Mayhew, 2013).





Larvae of first and second stage (freely moving) have a rhabditelike oesophagus with two bulbuses. Their intestinal cells are not distinguishable. The parasitic form (third stage) is distinguished by a filariform oesophagus with no extensions; the intestine does not have cells which can be clearly distinguished through an optical microscope (Fig. 1).

After cultivation, the liquid with the larvae was put in test tubes (10 ml) by 4 ml and was centrifuged for four minutes at 1500 revolutions a minute. 1 ml of sediments with larvae was evenly weighed and put in 0.1 ml portions in plastic containers with a capacity of 1.5 ml. After that, 1 ml of the substance under research was added to the larval culture (20–40 ind.), which then was kept for 24 hours at a temperature of 22–24 °C. Four concentrations (10, 1, 0.1 и 0.01 g/l) of each of the flavouring agents were used in the experiments (Table 1), with eight replications.

Results

The flavouring agents citral, benzaldehyde exhibited the strongest effect on the viability of *S. ransomi* larvae (Table 2).

Table 1Properties and usage of flavouring agents* which were used for establishing the viability level of *S. ransomi* larvae

Name of the substance	Chemical formula	Structural formula	Properties	Content	Usage	
					in food industry	in medicine
Benzaldehyde	C ₆ H ₅ CHO		colourless liquid with odour of bitter almond or of apple pips	cores of kernels of bitter almond, apricot kernels, stones of peaches, cherries, black cherries and other stone fruits, leaves of bird cherry tree, pulp of oyster mushroom (<i>Pleurotus ostreatus</i>)	as food additive	no data concerning usage
D-limonene (1-methyl-4-isopropenylcyclohexene -1)	C ₁₀ H ₁₆		colourless liquid with odour of citrus	many essential oils, including citrus	as food additive	no data concerning usage
Citral (3,7-dimethyl-2,6-octadienal)	C ₁₀ H ₁₆ O		viscous colourless or bright-yellow fluid with odour of lemon	essential oil of lemon grass (citronella), oil of cubeb, lemon essential oil, eucalyptus and some other essential oils	as food additive	antiseptics anti-inflammatory agent, used to relieve intracranial hypertension
Beta-ionone ((3E)-4-(2,6,6-trimethylcyclohex-1-enyl)but-3-en-2-one)	C ₁₃ H ₂₀ O		has a pleasant flower odour	some essential oils	as food additive	no data concerning usage

Note: * – Official Journal of the European Union, L 354/46 (2008), Sun (2007), Smirnov (2008).




Fig. 1. *S. ransomi* larvae: *a* – noninvasional stage, *b* – filariform oesophagus, *c* – caudal end, *d* – outlet from the cuticle: length of black bar – 50 μ m

Table 2
LD₅₀ (x ± SD) for *S. ransomi* larvae in laboratory experiment

Название вещества	LD ₅₀ , мг/л
Benzaldehyde	142 ± 64
D-limonene	–
Citral	97 ± 36
Beta-ionone	–

Note: «←» – the experiment did not achieve death of 50% larvae (needed concentration of more than 10 g/l).

D-limonene and beta-ionone at the studied concentrations (less than 10 g/m) did not affect the viability of *S. ransomi* larvae significantly, which shows that they are not useful as nematocidal medicines. The maximum effect upon the nematodes at LD₉₀ (Fig. 2a, c) was shown by benzaldehyde at 685 mg/l. In this case, the larvae of the 100% nematodes studied perished in benzaldehyde solutions with a concentration of 1 and 10 g/l. Thus, this substance can be used for further development of veterinary drugs with anthelmintic effect.

Discussion

Nowadays research is being conducted on the extent of effect of flavouring agents on the viability of agents of infection. Chiang et al. (2005) discovered the antiviral function of a broad spectrum of apigenin, linalool and ursolic acid. These substances are extracted from basil, which is familiar in Chinese medicine as a medicinal plant. The antimicrobial function of these compounds such as linalool and citral is well attested. They are capable of inhibiting the growth of pathogenic microorganisms (Sato et al., 2006; Somolinos

et al., 2008; Si et al., 2009; Belletti et al., 2010). Research on the effect of citral on infectious agents also supports the findings from our experiments on the viability of *S. ransomi* larvae. The effect of benzaldehyde and beta-ionone has been insufficiently studied.

For pest control in agriculture a number of authors have suggested using the food flavouring agent cinnamic aldehyde and also acaricidal substances (Knoblauch and Fry, 2011; Na et al., 2011; Shen et al., 2012; Belkind et al., 2013). According to Shen et al. (2012), after using this flavouring agent for 24 hours, LD₅₀ for *Psoroptes* was 107 mg/ml. Na et al. (2011) used it against *Dermatophagoides* of birds: LD₅₀ after 24 hours was 0.54 mg/ml. This substance was also tested by Cheng et al. (2009) against mosquito larvae: during 24 hours LD₅₀ was 40.8 mg/ml (LD₉₀ = 81.7 mg/ml) and 46.5 mg/ml (LC₉₀ = 83.3 mg/ml) for cinnamic aldehyde and cinnamic acetate, respectively. According to Lee et al. (2008), cinnamic aldehyde, benzaldehyde, linalool, limonene, alpha-Terpineol and other flavouring agents have insecticidal properties against *Stophilus oryzae* (Linnaeus, 1763) (Coleoptera, Curculionidae): LD₅₀ at 48 hours exposure was 0.004–0.200 mg/cm².

As a fungicidal substance it is advised to use E₂₁₀ (Codex Alimentarius), or benzoic acid, which is also included in other food additives, such as E₂₁₁ – Sodium Benzoate, E₂₁₂ – Potassium Benzoate, E₂₁₃ – Calcium dibenzoate (Beerse et al., 2001; Amborabe et al., 2002; Joshi et al., 2008).

Methylparabene is used against fungi, and also as an antiseptic (Shapiro et al., 2002; Posey et al., 2005; Kromidas et al., 2006; Rebbeck et al., 2006; Ishiwatari et al., 2007; Gopalakrishnan et al., 2012). This substance is also included as a preservative in an insecticide aimed at controlling agricultural pests (Bell, 1990).




Fig. 2. The effect of benzaldehyde (a), D-limonene (b), citral (c) and beta-ionone (d) on the viability of *S. ransomi* larvae: along the axes of abscissa – variant of experiment (mg/l), along the axes of ordinate – viability of *S. ransomi* larvae during 24 hours of laboratory experiment (%); n = 8 for each variant of experiment

The flavouring agent benzaldehyde also shows insecticidal properties. The negative effect of benzaldehyde upon *Galleria mellonella* (Linnaeus, 1758) has been demonstrated. Therefore it is recommended for use in developing new agricultural insecticidal preparations (Ullah et al., 2015). The effect of beta-ionone upon living organisms has received insufficient study. Indeed, the effect of entire groups of flavouring agents upon living organisms requires further study in general.

Conclusions

We researched the effect of flavouring agents such as benzaldehyde, citral, D-limonene and beta-ionone upon the viability of *S. ransomi*, parasitic larvae on pigs. We determined the minimum values of LD₅₀ for benzaldehyde and citral. Experimentally, a death rate of 50% of the tested larvae was not achieved using D-limonene and beta-ionone. Food additives with a pleasant flower odour, which are permitted to be used for food, and which are used as cosmetics, are important for evaluating potential new antihelminthic medical and veterinary preparations.

References

Amborabe, B.-E., Fleurat-Lessard, P., Chollet, J. F., & Roblin, G. (2002). Antifungal effects of salicylic acid and other benzoic acid derivatives towards *Eutypa lata*: Structure – activity relationship. *Plant Physiology and Biochemistry*, 40, 1051–1060.

Beerse, P. W., Biedermann, K. A., Page, S. H., Mobley, M. J., & Morgan, J. M. (2001). Antimicrobial compositions comprising a benzoic acid analog and a metal salt. U.S. Patent No. 6294186. Libertyville, IL, USA.

Belkind, B. A., Shammo, B., Dickenson, R., Rehberger, L. A., & Heiman, D. F. (2013). Cinnamaldehyde and diallyl disulfide formulations and methods of use. U.S. Patent No. 20130018107. Libertyville, IL, USA.

Bell, M. (1990). Insecticidal aqueous-based microemulsion compositions. U.S. Patent No. 4904695. Stamford, CT, USA.

Belletti, N., Kamdem, S. S., Tabanelli, G., Lanciotti, R., & Gardini, F. (2010). Modeling of combined effects of citral, linalool and β-pinene used against *Saccharomyces cerevisiae* in citrus-based beverages subjected to a mild heat treatment. *International Journal of Food Microbiology*, 136(3), 283–289.

Boyko, O. O., Zazharska, N. M., & Brygadyrenko, V. V. (2016). The influence of the extent of infestation by helminths upon changes in body weight of sheep in Ukraine. *Visnyk of Dnipropetrovsk University. Biology, Ecology*, 24(1), 3–7.

Burke, J. M., Wells, A., Casey, P., & Kaplan, R. M. (2009). Herbal dewormer fails to control gastrointestinal nematodes in goats. *Veterinary Parasitology*, 160(1–2), 168–170.

Charlier, J., Duchateau, L., Claerebout, E., Williams, D., & Vercruyse, J. (2007). Associations between anti-*Fasciola hepatica* antibody levels in bulk-tank milk samples and production parameters in dairy herds. *Preventive Veterinary Medicine*, 78(1), 57–66.

Cheng, S.-S., Liu, J.-Y., Huang, C.-G., Hsui, Y.-R., Chen, W.-J., & Chang, S.-T. (2009). Insecticidal activities of leaf essential oils from *Cinnamomum osmophloeum* against three mosquito species. *Bioresource Technology*, 100, 457–464.

Chiang, L.-C., Ng, L.-T., Cheng, P.-W., Chiang, W., & Lin, C.-C. (2005). Antiviral activities of extracts and selected pure constituents of *Ocimum basilicum*. *Clinical and Experimental Pharmacology and Physiology*, 32(10), 811–816.

Cringoli, G., Veneziano, V., Jackson, F., Vercruyse, J., Greer, A. W., Fedele, V., Mezzino, L., & Rinaldi, L. (2008). Effects of strategic anthelmintic treatments on the milk production of dairy sheep naturally infected by gastrointestinal strongyles. *Veterinary Parasitology*, 156(3–4), 340–345.

Eysker, M., Boerdam, G. A., Hollanders, W., & Verheijden, J. H. M. (1994). The prevalence of *Isospora suis* and *Strongyloides ransomi* in suckling piglets in the Netherlands. *Veterinary Quarterly*, 16(4), 203–205.

Faye, D., Leak, S., Noulala, S., Fall, A., Losson, B., & Geerts, S. (2003). Effects of gastrointestinal helminth infections and plane of nutrition on the health and productivity of F1 (West African Dwarf × Sahelian) goat crosses in The Gambia. *Small Ruminant Research*, 50(1–2), 153–161.

Gopalakrishnan, S., Chitra, T. A., Aruna, A., & Chenthihnanathan, A. (2012). Development of RP-HPLC method for the simultaneous estimation of ambroxol hydrochloride, cetirizine hydrochloride and antimicrobial preservatives in combined dosage form. *Scholars Research Library, Der Pharma Chemica*, 4(3), 1003–1015.

Ishiwatari, S., Suzuki, T., Hitomi, T., Yoshino, T., Matsukuma, S., & Tsuji, T. (2007). Effects of methyl paraben on skin keratinocytes. *Journal of Applied Toxicology*, 27, 1–9.

Joshi, S. D., Vagdevi, H. M., Vaidya, V. P., & Gadaginamath, G. S. (2008). Synthesis of new 4-pyrrol-1-yl benzoic acid hydrazide analogs and some derived oxadiazole, triazole and pyrrole ring systems: A novel class of potential antibacterial and antitubercular agents. *European Journal of Medicinal Chemistry*, 43, 1989–1996.

Knecht, D., Popiolek, M., & Zalesny, G. (2011). Does meatiness of pigs depend on the level of gastro-intestinal parasites infection? *Preventive Veterinary Medicine*, 99(2–4), 234–239.

Knoblauch, C., & Fry, K. (2011). Non-toxic insecticide. U.S. Patent No. 7956092. Edmonton, Alberta, CA, USA.

Kromidas, L., Perrier, E., Flanagan, J., Rivero, R., & Bonnet, I. (2006). Release of antimicrobial actives from microcapsules by the action of axillary bacteria. *International Journal of Cosmetic Science*, 28, 103–108.

Lee, E.-J., Kim, J.-R., Choi, D.-R., & Ahn, Y.-J. (2008). Toxicity of cassia and cinnamon oil compounds and cinnamaldehyde-related compounds to *Sitophilus oryzae* (Coleoptera: Curculionidae). *Journal of Economic Entomology*, 101(6), 1960–1966.

Lu, C. D., Gangyi, X., & Kawas, J. R. (2010). Organic goat production, processing and marketing: Opportunities, challenges and outlook. *Small Ruminant Research*, 89(2–3), 102–109.

Maslova, E. N., Sidorova, K. A., Antropov, V. A., & Dragich, O. A. (2015). Otsenka tyazhesti techeniya boleznii pri parazitozakh svinej [Assessment of the severity of the disease in parasitic diseases of pigs]. *Modern Problems of Science and Education*, 3, 1–8 (in Russian).

Na, Y. E., Kim, S.-I., Bang, H.-S., Kim, B.-S., & Ahn, Y.-J. (2011). Fumigant toxicity of cassia and cinnamon oils and cinnamaldehyde and structurally related compounds to *Dermatophyllum gallinae* (Acari: Dermatophyidae). *Veterinary Parasitology*, 178, 324–329.

Ponamar, S. I., Shendrik, L. I., Shendrik, C. M., Gugosyan, Y. A., & Shendrik, I. M. (2014a). Izmenenie gematologicheskikh pokazatelej do i posle lecheniya telyat pri strongiloidoze [Changes in hematology parameters before and after treatment of calves reached strongyloidiasis]. *Science and Technology Bulletin of Scientific Research Center for Biosafety and Environmental Control of Agro-Industrial Complex*, 2(2), 112–118 (in Ukrainian).

Ponamar, S. I., Goncharenko, V. P., Kruchinenko, O. V., & Shendrik, C. N. (2014b). Effektivnost' kompleksnogo podkhoda pri postanovke diagnoza na strongiloidoz [Effectiveness of the integrated approach in the diagnosis of strongyloidiasis]. *Scientific Messenger of Veterinary Medicine*, 13, 190–193 (in Ukrainian).

Ponamar, S. I., Soroka, N. M., & Ponamar, Z. S. (2013). Kompleksnaya ehtiotropnaya terapiya svinej pri nematodozakh [Complex ethiotropic therapy for pigs infected by nematodes]. *Veterinary*, 2, 29–31 (in Russian).

Posey, R., Culbertson, E. C., & Westermeier, J. C. (2005). Clear barrier coating and coated film. U.S. Patent No. 6911255. Greer, SC, USA.

Rahmann, G., & Seip, H. (2006). Alternative strategies to prevent and control endoparasite diseases in organic sheep and goat farming systems. *Ressortforschung für den Ökologischen Landbau*, 298, 49–90.

Rebbeck, C., Hammond, R., Wong, J., Nair, L., & Raghavan, N. (2006). Solid-phase extraction and HPLC analysis of methylparaben and propylparaben in a concentrated antibiotic suspension. *Drug Development and Industrial Pharmacy*, 32, 1095–1102.

Samsonovich, V. A. (2012a). Ehkologicheskie osobennosti ehpizootologii strongiloidoza porosyat [Ecological features of epizootology strongyloidosis pigs]. *Scientific Notes of the Bauman Kazan State Academy of Veterinary Medicine*, 211, 135–138 (in Russian).

Samsonovich, V. A. (2012b). Vliyanie strongiloidov na morfologicheskij sostav krovi i pokazateli estestvennoj rezistentnosti u porosyat [Influence of *Strongyloides* on the morphological composition of the blood and indicators of natural resistance in pigs]. *Proceedings of the National Academy of Sciences of Belarus. Series of Agrarian Sciences*, 3, 74–79 (in Russian).

Sato, K., Krist, S., & Buchbauer, G. (2006). Antimicrobial effect of trans-Cinnamaldehyde, (–)-Perillaldehyde, (–)-Citronellal, Citral, Eugenol and Carvacrol on airborne microbes using an airwasher. *Biological and Pharmaceutical Bulletin*, 29(11), 2292–2294.

Shapiro, S., Giertsen, E., Guggenheim, B. (2002). An *in vitro* oral biofilm model for comparing the efficacy of antimicrobial mouthrinses. *Caries Research*, 36, 93–100.

Shen, F., Xing, M., Liu, L., Tang, X., Wang, W., Wang, X., Wu, X., Wang, X., Wang, X., Wang, G., Zhang, J., Li, L., Zhang, J., & Yu, L. (2012). Efficacy of trans-cinnamaldehyde against *Psoroptes cuniculi* *in vitro*. *Parasitology Research*, 110, 1321–1326.

Si, W., Ni, X., Gong, J., Yu, H., Tsao, R., Han, Y., & Chambers, J. R. (2009). Antimicrobial activity of essential oils and structurally related synthetic food additives towards *Clostridium perfringens*. *Journal of Applied Microbiology*, 106(1), 213–220.

- Smirnov, E. V. (2008). Pishhevye aromatizatory [Food flavors]. Profession, Saint Petersburg (in Russian).
- Somolinos, M., Garcia, D., Pagan, R., & Mackey, B. (2008). Relationship between sublethal injury and microbial inactivation by the combination of high hydrostatic pressure and citral or tert-butyl hydroquinone. *Applied and Environmental Microbiology*, 74(24), 7570–7577.
- Sun, J. (2007). D-Limonene: Safety and clinical applications. *Alternative Medicine Review*, 12(3), 259–264.
- Ullah, I., Latif, A. K., Ali, L., Khan, A. R., Waqas, M., Hussain, J., Lee, I.-J., & Shin, J.-H. (2015). Benzaldehyde as an insecticidal, antimicrobial, and antioxidant compound produced by *Photobacterium temperata*. *Journal of Microbiology*, 53(2), 127–133.
- Van Wyk, J. A., & Mayhew, E. (2013). Morphological identification of parasitic nematode infective larvae of small ruminants and cattle: A practical lab guide. *Onderstepoort Journal of Veterinary Research*, 80(1), 1–14.
- Veneziano, V., Rubino, R., Fedele, V., Rinaldi, L., Santaniello, M., Schioppi, M., Cascone, C., Pizzillo, M., & Cringoli, G. (2004). The effects of five anthelmintic treatment regimes on milk production in goats naturally infected by gastrointestinal nematodes. *South African Journal of Animal Science*, 34, 238–240.
- Yamov, V. Z., & Antropov, V. A. (2008). Ehpizootologiya gel'mintozov svinej na tyumenskom yuge [Epizootology of pigs helminthosis in the south of Tyumen]. *Agricultural Messenger of Ural Veterinary Science*, 47, 70–71 (in Russian).
- Zajac, A. M., & Conboy, G. A. (eds.) (2011). *Veterinary clinical parasitology*, 8th ed. John Wiley and Sons, UK.